

ApJ; Submitted Dec 13, 2000; accepted Jan 30, 2001

Magnetically Accreting Isolated Old Neutron Stars

Robert E. Rutledge¹

ABSTRACT

Previous work on the emission from isolated old neutron stars (IONS) accreting the inter-stellar medium (ISM) focussed on gravitational capture – Bondi accretion. We propose a new class of sources which accrete via magnetic interaction with the ISM. While for the Bondi mechanism, the accretion rate $\dot{M}_{\text{Bondi}}$ decreases with increasing NS velocity, in magnetic accretors (MAGACs) $\dot{M}_{\text{MAGAC}}$ *increases* with increasing NS velocity ($\dot{M}_{\text{Bondi}} \propto v^{-3}$ vs. $\dot{M}_{\text{MAGAC}} \propto v^{1/3}$). MAGACs will be produced among high velocity ($\gtrsim 100 \text{ km s}^{-1}$) high magnetic field ($B > 10^{14} \text{ G}$) radio pulsars – the “magnetars” – after they have evolved first through magnetic dipole spin-down, followed by a “propeller” phase (when the object sheds angular momentum on a timescale $\lesssim 10^{10} \text{ yr}$). The properties of MAGACs may be summarized thus: dipole magnetic fields of $B \gtrsim 10^{14} \text{ G}$; minimum velocities relative to the ISM of $> 25\text{--}100 \text{ km s}^{-1}$, depending on B , well below the median in the observed radio-pulsar population; spin-periods of $>$ days to years; accretion luminosities of $10^{28}\text{--}10^{31} \text{ erg s}^{-1}$; and effective temperatures $kT_{\text{eff}} = 0.3\text{--}2.5 \text{ keV}$ if they accrete onto the magnetic polar cap. We find no examples of MAGACs among previously observed source classes (anomalous X-ray pulsars, soft-gamma-ray repeaters or known IONS). However, MAGACs may be more prevalent in flux-limited X-ray catalogs than their gravitationally accreting counterparts.

Subject headings: pulsars: general – stars: magnetic fields – stars: neutron – X-rays: stars

¹ Space Radiation Laboratory, California Institute of Technology, MS 220-47, Pasadena, CA 91125; rutledge@srl.caltech.edu; and Institute for Theoretical Physics, Kohn Hall, University of California, Santa Barbara, CA 93106

1. Introduction

Accretion onto isolated old neutron stars (IONS) through the spherical, Bondi-accretion mode (Bondi & Hoyle 1944; Bondi 1952) has been investigated as a means to produce a population of neutron stars detectable in the ROSAT X-ray All Sky Survey (RASS; Treves & Colpi 1991; Blaes & Madau 1993; Madau & Blaes 1994; Zane et al. 1995; Popov et al. 2000). The estimated number of these sources in the RASS was initially high ($\sim 10^3$ – 10^4). However, as observations of higher mean velocities in the radio pulsar population (Lorimer et al. 1997; Hansen & Phinney 1997; Cordes & Chernoff 1998, and references therein) were considered, the predicted number of detectable IONS accreting from the ISM decreased dramatically ($\sim 10^2$ – 10^3). This is because the Bondi accretion rate is a strong function of the NS velocity through the interstellar medium (ISM)² –

$$\dot{M}_{\text{Bondi}} = 4\pi(GM_{\text{NS}})^2\rho v^{-3} \quad (1)$$

where ρ is the local mass density of the ISM. Bondi-accreting IONS are expected to have X-ray luminosities $\sim 10^{31}$ erg s^{−1} and effective temperatures $kT_{\text{eff}} \lesssim 200$ eV, both dependent upon the accretion rate.

There will be a different accretion mode, however, when the NS magnetic field dominates over the gravitational field in attracting accretion from the ISM. In this mode, the warm ionized medium (WIM) attaches to the NS magnetic field at the magnetosphere, accreting directly onto the NS surface. These magnetically accreting isolated old neutron stars (MAGACs, “magics”) contrast with IONS which accrete through the Bondi-mode, in that the Bondi accretion rate decreases with the NS velocity, while in MAGACs, the accretion rate *increases* with increasing velocity:

$$\dot{M}_{\text{MAGAC}} = \pi R_{\text{M}}^2 v \rho \quad (2)$$

where $R_{\text{M}} = (\mu^2/(4\pi\rho v^2))^{1/6}$ is the size-scale for the magnetosphere of a non-rotating NS, where the magnetic energy density from the NS’s magnetic moment (μ) is equal to the ram-pressure of the WIM. This gives $\dot{M}_{\text{MAGAC}} \propto v^{1/3}$. So, for a given magnetic field strength, Bondi accretion dominates at low velocities, while magnetic accretion dominates at high velocities.

²Here, $v = \sqrt{V^2 + C_s^2}$, the geometric sum of the spatial velocity of the NS and the sound speed in the accreted medium. The sound speed of the ISM is low (~ 10 km s^{−1}) compared to spatial velocities relevant to the present work ($v \gtrsim 100$ km s^{−1}), so we assume $v \gg C_s$ throughout.

The evolutionary scenario of magnetically accreting NSs has been outlined previously, by Harding & Leventhal (1992, HL92 hereafter). HL92 found magnetised NSs first spin down via dipole radiation, followed by a longer spin-down period via the propellor effect. They restricted their attention to sources with surface magnetic fields $\lesssim 10^{13}$ G, and found that they are unable to accrete onto the compact object, as the propellor spin-down time is longer than the age of the universe. A similar line of argument is followed in the review by Treves et al. (2000) in the context of Bondi accreting IONS. This limitation does not apply to NSs with magnetic fields $\gtrsim 10^{14}$ G; a NS with magnetic field $B = 10^{15}$ G and velocity $v = 300$ km s $^{-1}$, spins down in $\sim 4 \times 10^9$ yr.

On the other hand, this same evolutionary scenario was used to argue for magnetic field decay in the IONS RX J0720.4-3125 (Wang 1997; Konenkov & Popov 1997), and for NSs in general (Livio et al. 1998; Colpi et al. 1998). We suggest, instead, that high magnetic field IONS have not been discovered due to the fact that strong magnetic field sources accreting from the ISM are spectrally harder (we find below) than Bondi accretors, and have been selected against by observers, or confused with background AGN.

Moreover, since HL92’s work, observations have revealed NSs with considerably stronger magnetic fields than considered by HL92, in the range of 10^{15} G – the magnetars. Pulse-timing of soft-gamma-ray repeaters (Kouveliotou et al. 1998; Kouveliotou et al. 1999) confirms their existence as proposed in earlier theoretical work (Thompson & Duncan 1993; Duncan & Thompson 1992). Circumstantial evidence suggests that anomalous X-ray pulsars have similarly strong B -fields (Thompson & Duncan 1996; Vasisht & Gotthelf 1997; Heyl & Hernquist 1997); and precision pulsar-timing evidence also implies magnetic fields in excess of 10^{14} G (Kaspi et al. 1999), confirming less-precise multi-epoch timing solutions which nonetheless led to a magnetar interpretation (Gotthelf et al. 1999). Static solutions for NSs have been proposed for surface fields up to $\sim 10^{18}$ G (Bocquet et al. 1995; Cardall et al. 2001), although no objects with magnetic fields as high as this have yet been observed. Thus, with theoretical and observational motivation, we reconsider the scenario of HL92, but with particular attention paid to NSs with magnetic fields above 10^{13} G.

In §2 we describe the model for this population, following the scenario of HL92. In §3, we summarize the properties of MAGACs, discuss observational ramifications and conclude.

2. Model

2.1. Evolutionary Model

We assume that the neutron star is born with a velocity $v = v_7 10^7 \text{ cm s}^{-1}$ relative to the inter-stellar medium, and a magnetic moment $\mu = \mu_{33} 10^{33} \text{ G cm}^3$, which corresponds to a dipole field strength at the NS surface (10 km) of 10^{15} G (appropriate for magnetars; Thompson & Duncan 1993; Duncan & Thompson 1992). The spin period is initially short ($P \ll 10 \text{ sec}$). We assume the magnetic field does not decay. We assume a compact object mass $M = M_{\text{NS}} 1.4 M_{\odot}$, radius $R = R_{\text{NS}} 10^6 \text{ cm}$. We assume the NS travels through a spatially uniform proton density $n_p = 0.1 n_{-1} \text{ cm}^{-3}$ in the WIM (Boulares & Cox 1990; Ferriere 1998a; Ferriere 1998b). A more detailed examination of the properties of this population, which would include a distribution of NS kick velocities, orbits about a realistic galactic potential, and the galactic gas density and ionization state – such as that performed for Bondi accretors and cooling NSs (Popov et al. 2000; Popov 2001) – will be the subject of forthcoming work.

First, HL92 argues that magnetic accretion dominates over Bondi accretion when the magnetospheric radius $R_{\text{M}} = 4.2 \times 10^{12} \mu_{33}^{1/3} v_7^{-1/3} n_{-1}^{-1/6} \text{ cm}$ is larger than the Bondi accretion radius $R_{\text{Bondi}} = \frac{2GM_{\text{NS}}}{v^2}$. This inequality results in a velocity limitation for a MAGAC:

$$v_{\text{lim}} > 94 \mu_{33}^{-1/5} n_{-1}^{1/10} M_{\text{NS}}^{3/5} \text{ km s}^{-1} \quad (3)$$

Electromagnetic dipole radiation pressure inhibits accretion from the ISM until the light cylinder is external to the magnetospheric radius ($R_{\text{L.C.}} > R_{\text{M}}$), which occurs at a spin period of

$$P_{\text{L.C.}} = 880 \mu_{33}^{1/3} v_7^{-1/3} n_{-1}^{-1/6} \text{ s} \quad (4)$$

The timescale to reach this period through dipole radiation is:

$$\tau_{\text{L.C.}} = 2.3 \times 10^7 I_{45} \mu_{33}^{-4/3} v_7^{-2/3} n_{-1}^{-1/3} \text{ yr} \quad (5)$$

where I_{45} is the moment of inertia in units of 10^{45} g cm^2 . This is a short timescale, compared with the life of the NS. After reaching this period, the light cylinder is at the R_{M} ; however, the NS still cannot accrete (or, accretes inefficiently) due to the “propeller effect” (Illarionov & Sunyaev 1975) in which matter is spun away by the magnetosphere, which is outside the Keplerian orbit in co-rotation with the neutron star (the co-rotation radius). This matter is ejected from the system, carrying angular momentum with it, spinning down the NS until the co-rotation radius reaches R_{M} , at which point the NS will have a spin period:

$$P_P = 3.9 \times 10^6 \mu_{33}^{1/2} v_7^{-1/2} n_{-1}^{-1/4} M_{\text{NS}}^{-1/2} \text{ s} \quad (6)$$

This spin period is substantially longer (by three orders of magnitude) than any spin period yet observed from a NS. The propeller spin-down timescale, when the accretion rate is dominated by magnetospheric accretion, is (Illarionov & Sunyaev 1975; Davies et al. 1979):

$$\tau_P = 2.3 \times 10^{10} I_{45} \mu_{33}^{-4/3} v_7^{-5/3} n_{-1}^{-1/3} \text{ yr} \quad (7)$$

The τ_P timescale dominates the evolutionary time between NS birth and emergence of the NS as a magnetic accretor. It is comparable to the age of the Galaxy ($\lesssim 10^{10}$ yr) velocities already observed among the NS population. but only for high-magnetic field sources ($\gtrsim 10^{15}$ G). This makes magnetic accretors with the right combination of magnetic field and velocity observable.

Following spin-down via the propeller, matter is accreted onto onto the polar cap of the NS. The maximum accretion rate onto the compact object is (HL92):

$$\dot{M}_{\text{MAGAC}} = \pi R_M^2 v \rho = 9.6 \times 10^7 \mu_{33}^{2/3} v_7^{1/3} n_{-1}^{2/3} \text{ g s}^{-1} \quad (8)$$

This $\dot{M}_{\text{MAGAC}}$ is an upper-limit to the mass accretion rate onto the NS surface, and is potentially much lower (see §2.2). Moreover, the Bondi and MAGAC accretion rates are simply related to the magnetospheric and Bondi radius:

$$\frac{\dot{M}_{\text{MAGAC}}}{\dot{M}_{\text{Bondi}}} = \left(\frac{R_M}{R_{\text{Bondi}}} \right)^2 \quad (9)$$

If we assume free-fall accretion onto the NS surface, this provides an accretion luminosity of:

$$L_{\text{MAGAC}} = 1.8 \times 10^{28} \mu_{33}^{2/3} v_7^{1/3} n_{-1}^{2/3} M_{\text{NS}} R_{\text{NS}}^{-1} \text{ erg s}^{-1}. \quad (10)$$

To estimate the effective temperature, we take the size of the polar cap to be $R_{\text{cap}} = R_{\text{NS}}(R_{\text{NS}}/R_M)^{1/2}$. This would seem unusually small – ~ 50 m for $B = 10^{15}$ G. Nonetheless, the effective temperature T_{eff} of this emission is:

$$kT_{\text{eff}} = 0.39 \mu_{33}^{1/4} n_{-1}^{1/8} M_{\text{NS}}^{1/4} R_{\text{NS}}^{-1} \text{ keV}. \quad (11)$$

Note that kT_{eff} is independent of the NS velocity. The accretion luminosity will remain below the local Eddington luminosity if:

$$\frac{\sigma_{\text{es}}}{m_H} \frac{L_{\text{MAGAC}}}{c \pi R_{\text{cap}}^2} \left(\frac{R_{\text{NS}}}{r} \right)^2 < \frac{GM_{\text{NS}}}{r^2} \quad (12)$$

where m_H is the proton mass. The electron scattering cross-section in a strong magnetic field σ_{es} is suppressed from the Thomson rate σ_T by $\sigma_{\text{es}}/\sigma_T = 4 \times 10^{-6} (E/1 \text{ keV})^2 (B_{\text{QED}}/B)^2$, with $B_{\text{QED}} = 4.4 \times 10^{14} \text{ G}$, and photon energy E (Herold 1979). This result gives the constraint $\mu_{33}^{-1} n_{-1}^{1/2} < 3 \times 10^9$. This will not be violated for NSs with $B > 10^{13} \text{ G}$ unless the ISM is un-physically over-dense ($n_{-1} \gtrsim 10^6$). Thus, the accretion rate will be locally sub-Eddington.

Finally, the ratio of $\dot{M}_{\text{MAGAC}}$ to $\dot{M}_{\text{Bondi}}$ is:

$$\frac{\dot{M}_{\text{MAGAC}}}{\dot{M}_{\text{Bondi}}} = 1.8 \mu_{33}^{2/3} v_7^{10/3} n_{-1}^{-1/3} M_{\text{NS}}^{-2} \quad (13)$$

Thus, for the same velocity distribution, MAGACs will be more luminous than Bondi accretors. We discuss the observational ramifications of this in §3.

2.2. Comments on the Accretion Luminosity

We have two comments on the accretion luminosity. The first regards the accretion at magnetopause, and the second regards the need for the ISM to be ionized in order to be accreted.

Arons & Lea (1976) performed detailed analytic description of mass transfer across magnetopause in a 10^{12} G NS system, fed by a $\approx 600 \text{ km s}^{-1}$ wind from its companion, but with a substantially higher accretion rate at magnetopause ($6 \times 10^{19} \text{ g s}^{-1}$) than we consider here. Even at this lower magnetic field strength and higher mass accretion rate, they find a cusp forms with a stand-off shock, causing an interchange instability which mediates accretion onto the compact object. While inflow at the cusp increases with increasing density, the interchange instability operates more efficiently and prevents very high densities from arising at magnetopause. Three-dimensional MHD simulations find similar results (Toropin et al. 1999).

If the interchange instability of Arons and Lea (1976) permits accretion across the entire NS surface (rather than just at the magnetic poles), then the luminosity would remain the same, while dramatically decreasing the effective temperature, to:

$$kT_{\text{eff}} = 0.0076 \mu_{33}^{1/3} v_7^{1/12} n_{-1}^{1/6} M_{\text{NS}}^{1/4} R_{\text{NS}}^{-3/4} \text{ keV}. \quad (14)$$

Further analysis of accretion onto a magnetic dipole with strength 10^{13} – 10^{18} G is necessary to estimate the magnitude of the effect on the luminosity and spectrum. For now, we adopt $\dot{M}_{\text{MAGAC}}$ as an upper-limit.

Secondly, to accrete material via magnetic interaction, the material must be ionized. Throughout, we have therefore assumed that MAGACs accrete from the (ionized) WIM, and not from the neutral ISM. However, one might wonder if the MAGAC’s luminosity is sufficient to ionize the ISM by the time it reaches the magnetosphere (assuming that accretion is already taking place). Ionization of the ISM by an accreting IONS has been examined previously (Shvartsman 1971; Blaes et al. 1995). These investigations found that a Bondi-accretion luminosity was capable of ionizing the ISM well beyond the Bondi-accretion radius. However, we have applied a similar approach to MAGACs accreting onto the magnetic polar cap, and find the fraction of a neutral ISM which would be ionized by the MAGAC at R_M to be $\ll 1$, for a broad range of velocities ($0.1 < v_7 < 10$) and densities ($0.1 < n_{-1} < 100$) for magnetic field strengths considered here. Therefore, MAGACs must accrete from the WIM, which has a typical density of $\sim 0.13 \text{ cm}^{-3}$ for a filling factor of 20% in the plane and a scale height of 1 kpc (Ferriere 1998b). This justifies our assumption that MAGACs must accrete from the WIM, and not from the neutral ISM.

2.3. Limits on the Observability of MAGACs

Here, we list those effects which will limit the observability of MAGACs, and which determine the luminosity, spectral hardness, velocity, magnetic field, and pulse-period parameter space in which they may be observable. In Fig. 1, we divide the NS B -field dipole strength, velocity parameter space, on the basis of these limits:

1. $\dot{M}_{\text{MAGAC}} > \dot{M}_{\text{Bondi}}$. As we find above, this results in a velocity limit (Eq. 3), above which the NS is a MAGAC, and below which the NS is a Bondi accretor (broken line).
2. *The propeller phase must be shorter than the age of the galaxy.* We use as a limit (Eq. 7) that $\tau_p < 1 \times 10^{10}$ yr (dotted line), and also include the line for $\tau_p < 1 \times 10^9$ yr. This age limit also effectively sets a magnetic field strength limit: NSs with $B < 10^{13}$ G will not spin down in a Hubble time. Thus, if magnetic fields in NSs decay to $< 10^{13}$ G in less than a Hubble time, they will not be observed as MAGACs.

3. *The pulsar must not escape the Galaxy.* We adopt an escape speed limit of 700 km s^{-1} (cf. Miyamoto & Nagai 1975; Paczynski 1990), although the exact value depends on where in the galaxy the NS is formed. This limit is indicated by the hashed area at $v > 700 \text{ km s}^{-1}$.
4. *The NS magnetic field should not exceed the stability limit.* We assume a surface dipole field of $B < 10^{18} \text{ G}$ (Bocquet et al. 1995; Cardall et al. 2001 ; dashed-dot line).

We see in Fig. 1 that acceptable magnetic fields above 10^{14} G , and acceptable velocities range from $\sim 25 \text{ km s}^{-1}$ to the escape speed from the galaxy. We include lines of constant L_{MAGAC} (for density = $0.1 \text{ proton cm}^{-3}$; Eq. 10); the luminosities of MAGACs are between 10^{28} and $10^{31} \text{ erg s}^{-1}$ (again, assuming free-fall accretion from the magnetosphere onto the NS). On the right axis of the figure, we mark kT_{eff} , which is independent of velocity, and only weakly dependent upon the density of the ISM ($\propto n^{1/8}$). MAGACs have $kT_{\text{eff}} = 0.3\text{--}2.5 \text{ keV}$. These are harder than observed from any of the identified IONSs to date (Treves et al. 2000), perhaps due to an observational bias.

Finally, we note that for values of $B = 10^{12} \text{ G}$ and $v = 400 \text{ km s}^{-1}$ which are not unusual in pulsars, such an object will not accrete via the Bondi-mode from the ISM, as the magnetic field is in the propeller range. In addition the source will not spin-down in $< 10^{10} \text{ yr}$ timescale; so it will wander the Galaxy as a dipole-spun down, propelling NS.

3. Discussion and Conclusions

Strong magnetic field ($B > 10^{14} \text{ G}$) NSs go through three stages of evolution: (1) dipole emission; (2) propeller slowdown; and finally, the MAGAC stage (3) magnetic accretion onto the NS surface. MAGACs have the following properties: surface magnetic fields between 10^{14} G up through the NS stability limit ($\sim 10^{18} \text{ G}$); spin periods between $10^6\text{--}10^8 \text{ s}$; X-ray luminosities $10^{28}\text{--}10^{31} \text{ erg s}^{-1}$, for a WIM density of $0.1 \text{ protons cm}^{-3}$; and effective temperatures $kT_{\text{eff}} = 0.3\text{--}2.5 \text{ keV}$. The low X-ray luminosities of these sources make it unlikely that the very long spin periods will be detected as pulsations with present X-ray instrumentation, as prohibitively long integrations with frequent monitoring would be required.

At the median velocity for radio pulsars, MAGACs will be $\sim \times 130$ more luminous than Bondi accretors (for $v_7 = 3$, a typical median velocity for pulsars; Lorimer et al. 1997; Hansen & Phinney 1997; Cordes & Chernoff 1998, and references therein). Although the birth-rate of magnetars has been estimated to be $\sim 10\%$ of isolated pulsars (Kouveliotou & et al. 1994; Kouveliotou et al. 1998), they can be observed to ≈ 10 times the distance

and $\approx 130\times$ the volume of the Galactic disk of Bondi accretors at the radio pulsar median velocity. Magnetic accretors, therefore, may be more prevalent in flux limited X-ray catalogs, such as the ROSAT All-Sky-Survey/Bright Source Catalog (RASS/BSC ; Voges et al. 1999) than the lower-magnetic field Bondi accretors. Note, however, that Popov et al. (Popov et al. 2000) found that the detected RASS/BSC INSs are more naturally interpreted as soft, young, nearby cooling neutron stars, rather than Bondi accretors; in this interpretation, no NSs accreting from the ISM have yet been discovered. The relative number of detectable MAGACs vs. Bondi accretors will ultimately depend on the particulars of the distribution of magnetic fields at birth in NSs, the kick velocities (and their possible dependence on B), as well as magnetic field decay, which we have neglected herein.

A brief comparison between observations and the predicted properties of this population reveal no strong candidates for MAGACs among known source classes.

Among radio pulsars, there are no known objects with implied B and v which would place them in the MAGAC B, v parameter space. The strongest B field radio pulsar is 5.5×10^{13} G (Camilo et al. 2000), is just below the low- B limit; its velocity is unmeasured. At a velocity of ~ 300 km s $^{-1}$, this NS will spin-down on a timescale of $\tau_P \approx 210^{11}$ yr, much longer than the age of the Galaxy. Thus, there are no observed radio pulsars in which the magnetic field is clearly strong enough to evolve into a MAGAC. This is possibly due to the relatively short timescale that such objects are dipole emitters ($\tau_{L.C.} \lesssim 10^6$ yr).

The observed spin periods of magnetars are $P \sim 1\text{--}15$ sec, which, for the magnetic fields implied (10^{15} G), are still within the range where electromagnetic dipole radiation will inhibit accretion. In addition, their X-ray luminosities are a factor $\gtrsim 1000$ larger than those we expect for MAGACs. These objects therefore are not presently MAGACs; however, depending on their spatial velocities, they may evolve into MAGACs.

The (model dependent) age of the isolated NS RX J185635-3754 (~ 1 My; Walter et al. 1996; Walter 2001) is shorter than τ_P , unless the NS has close to $B \sim 210^{18}$ G. With such a high magnetic field, along with the implied velocity (≈ 200 km s $^{-1}$), the source would be within the MAGAC parameter space. Its X-ray luminosity is $\approx 2\times 10^{31}$ erg s $^{-1}$, which is within a factor of a few of the expected luminosity for a MAGAC with this B and v . However, the effective temperature ($kT_{\text{eff}} = 0.057$ keV) is substantially below that expected from a MAGAC with $B \times 10^{18}$ G (2.6 keV; Eq. 11). The spectral hardness seems to rule out this object as a MAGAC for the polar-cap accreting scenario. Since all previously observed INSs have effective temperatures < 200 eV (Treves et al. 2000), it seems similarly unlikely that any known INS is a MAGAC. It is possible, however, that an interchange instability at the magnetosphere permits accretion over a larger fraction of the NS, which results in a softer spectrum, such as observed here (ref. 14).

Discovering these objects in the RASS/BSC requires a similar approach taken in previous work to identify IONS – identifying an X-ray source with no apparent optical counterpart, down to an X-ray-to-optical flux ratio of $f_x/f_{\text{opt}} \gtrsim 1000$, with confirmation coming from an optical detection at an appropriate value. However, MAGACs can be spectrally harder than their Bondi accreting counterparts ($kT_{\text{eff}} \sim 1 - 3$ keV; cf. Popov 2001), and searches for these objects must take this into account. Also, given their strong magnetic fields, these objects may have a cyclotron *emission* line, which may aid identification through X-ray spectroscopy (Nelson et al. 1995). Moreover, because we have assumed no magnetic field decay, the absence of MAGACs in the Galaxy, in combination with a population synthesis with high magnetic field SGRs and AXPs, would support field decay models.

We conclude that MAGACs may present a population which are equally prevalent in flux limited X-ray catalogs as Bondi accreting sources. However, no observed examples of this class are yet known.

I thank Lars Bildsten, Omer Blaes, David Chernoff and Chris Thompson for useful discussions. I thank Lars Bildsten and Omer Blaes for comments on the draft of this paper, which improved the manuscript. I am grateful to the participants in the ITP Workshop, “Spin and Magnetism in Young Neutron Stars” for stimulating discussions which brought about this work. This research was supported in part by the National Science Foundation under Grant No. PHY99-07949.

REFERENCES

- Arons, J. & Lea, S. M., 1976, ApJ 207, 914
- Blaes, O. & Madau, P., 1993, ApJ 403, 690
- Blaes, O., Warren, O., & Madau, P., 1995, ApJ 454, 370
- Bocquet, M., Bonazzola, S., Gourgoulhon, E., & Novak, J., 1995, A&A 301, 757
- Bondi, H., 1952, MNRAS 112, 195
- Bondi, H. & Hoyle, F., 1944, MNRAS 104, 421
- Boulares, A. & Cox, D. P., 1990, ApJ 365, 544
- Camilo, F., Kaspi, V. M., Lyne, A. G., Manchester, R. N., Bell, J. F., D’Amico, N., McKay, N. P. F., & Crawford, F., 2000, ApJ 541, 367

- Cardall, C., Prakash, M., & Lattimer, J., 2001, ApJ, submitted, astro-ph/0011148
- Colpi, M., Turolla, R., Zane, S., & Treves, A., 1998, ApJ 501, 252
- Cordes, J. M. & Chernoff, D. F., 1998, ApJ 505, 315
- Davies, R. E., Fabian, A. C., & Pringle, J. E., 1979, MNRAS 186, 779
- Duncan, R. C. & Thompson, C., 1992, ApJ 392, L9
- Ferriere, K., 1998a, ApJ 497, 759
- Ferriere, K., 1998b, ApJ 503, 700
- Gotthelf, E. V., Vasisht, G., & Dotani, T., 1999, ApJ 522, L49
- Hansen, B. M. S. & Phinney, E. S., 1997, MNRAS 291, 569
- Harding, A. K. & Leventhal, M., 1992, Nature 357, 388
- Herold, H., 1979, Phys. Rev. D 19, 2868
- Heyl, J. S. & Hernquist, L., 1997, ApJ 489, L67
- Illarionov, A. F. & Sunyaev, R. A., 1975, A&A 39, 185
- Kaspi, V. M., Chakrabarty, D., & Steinberger, J., 1999, ApJ 525, L33
- Konenkov, D. Y. & Popov, S. B., 1997, *AZh Letters* 23, 498
- Kouveliotou, C., Dieters, S., Strohmayer, T., van Paradijs, J., Fishman, G. J., Meegan, C. A., Hurley, K., Kommers, J., Smith, I., Frail, D., & Murakami, T., 1998, Nature 393, 235
- Kouveliotou, C. & et al., 1994, in G. Fishman, J. Brainered, & K. Hurley (eds.), *Gamma Ray Bursts: Second Workshop, Huntsville, AL 1993*, No. 307 in AIP Conference Proceedings, p. 167, American Institute of Physics
- Kouveliotou, C., Strohmayer, T., Hurley, K., van Paradijs, J., Finger, M. H., Dieters, S., Woods, P., Thompson, C., & Duncan, R. C., 1999, ApJ 510, L115
- Livio, M., Xu, C., & Frank, J., 1998, ApJ 492, 298+
- Lorimer, D. R., Bailes, M., & Harrison, P. A., 1997, MNRAS 289, 592
- Madau, P. & Blaes, O., 1994, ApJ 423, 748

- Miyamoto, M. & Nagai, R., 1975, PASJ 27, 533
- Nelson, R. W., Wang, J. C. L., Salpeter, E. E., & Wasserman, I., 1995, ApJ 438, L99
- Paczynski, B., 1990, ApJ 348, 485
- Popov, S. B., 2001, in *"Astrophysical Sources of High Energy Particles & Radiation"*, Erice, Italy November 2000., astro-ph/0101031
- Popov, S. B., Colpi, M., Treves, A., Turolla, R., Lipunov, V. M., & Prokhorov, M. E., 2000, ApJ 530, 896
- Shvartsman, V. F., 1971, *Sov. Astr.* 14
- Thompson, C. & Duncan, R. C., 1993, ApJ 408, 194
- Thompson, C. & Duncan, R. C., 1996, ApJ 473, 322
- Toropin, Y. M., Toropina, O. D., Savelyev, V. V., Romanova, M. M., Chechetkin, V. M., & Lovelace, R. V. E., 1999, ApJ 517, 906
- Treves, A. & Colpi, M., 1991, A&A 241, 107
- Treves, A., Turolla, R., Zane, S., & Colpi, M., 2000, PASP 112, 297
- Vasisht, G. & Gotthelf, E. V., 1997, ApJ 486, L129
- Voges, W., Aschenbach, B., Boller, T., Bräuninger, H., Briel, U., Burkert, W., Dennerl, K., Englhauser, J., Gruber, R., Haberl, F., Hartner, G., Hasinger, G., Kürster, M., Pfeiffermann, E., Pietsch, W., Predehl, P., Rosso, C., Schmitt, J. H. M. M., Truemper, J., & Zimmermann, H. U., 1999, A&A 349, 389
- Walter, F. M., 2001, ApJ, accepted, astro-ph/0009031
- Walter, F. M., Wolk, S. J., & Neuhauser, R., 1996, Nature 379, 233
- Wang, J. C. L., 1997, ApJ 486, L119
- Zane, S., Turolla, R., Zampieri, L., Colpi, M., & Treves, A., 1995, ApJ 451, 739

Fig. 1.— Magnetic field strength (B) vs. velocity for pulsars, showing regions where MAGAC IONS can be observed, and their accretion luminosities. The observability of MAGACs are limited at low velocities (left – dashed line) by a large magnetosphere (these then accrete always via Bondi); at high velocities (right – hashed area) by the likelihood they will escape the Galaxy’s gravitational field; at low B (below – dotted line) from the fact that the NSs will not spin down via the propeller mechanism in <10 Gyr (we also include a line for $\tau_P = 1$ Gyr; and at high B (above – dashed-dot line) by the instability of high B neutron stars. Note also that a moderate magnetic field NS (10^{12} G) with an average velocity (500 km s^{-1}) will not accrete gravitationally due to the propeller mechanism. We include lines of the maximum accretion luminosity for the MAGACs after being spun down via the propeller. The right axis shows the effective temperature for accretion onto the magnetic cap, which is independent of velocity and only weakly dependent on density ($\propto n^{1/8}$).




Figure 1